



# A Novel Generalized Divergence Measure of Fuzzy Sets

**Anshu Ohlan**

Assistant Professor, Department of Mathematics, All India Jat Heroes Memorial College, Rohtak Haryana, India.

## ABSTRACT

**Aim:** The aim of the paper is to introduce a generalized measure of discrimination in fuzzy environment.

**Methodology:** To achieve the goal of this paper a parametric generalization of Hellinger's fuzzy divergence measure is studied along with the proof of its validity.

**Results:** A particular case and some important properties are discussed in detail of the proposed generalized Hellinger's fuzzy divergence measure.

**Conclusion:** Generalized Hellinger's fuzzy divergence measure is valid measure of fuzzy divergence.

**Key Words:** Fuzzy sets, Fuzzy entropy, Fuzzy divergence measure, Generalized Fuzzy Divergence Measures

## INTRODUCTION

Shannon [1] was first to use the word "entropy" to measure an uncertain degree of the randomness in a probability distribution. The theory of fuzzy sets and fuzzy logic developed by Zadeh [2] has been used to form ambiguity, lack of information and uncertainty inherent in the human decision making process. It has achieved a great success in various areas such as multi-criteria decision making (MCDM), logical programming, pattern recognition, medical diagnosis and so on. Zadeh [3] introduced fuzzy entropy as an significant concept for measuring fuzzy information.

Thereafter, a number of researchers made study on the fuzzy theory and find their applications in different areas. For example, some new information and divergence measures and their applications in different areas have been proposed in literature [4-22].

Although many information measures between fuzzy sets has been emerging in the last decades. Still, there is a need to define quantitative information measures for imprecision, discrimination, distance, etc. over fuzzy sets with their practical applications. In literature the Hellinger's measure of discrimination was firstly introduced by Hellinger [23]. Here we propose a new parametric generalized Hellinger's divergence measure in fuzzy environment which provides a flexible approach to further leverage of choice to the user. It

may be observed that the potential, strength and efficiency of this new generalized Hellinger's fuzzy divergence measure exist in its properties.

Methodology section we recall and discuss some well-known concepts and the notions related to fuzzy set theory. In results section we introduce a generalized Hellinger's fuzzy divergence measure with the proof of validity. Some interesting properties of the proposed measure between different fuzzy sets are studied drawn in discussion section. Final section concludes the paper.

## METHODOLOGY

We now review a number of well-known concepts and definitions related with the theory of fuzzy sets. The uncertainty and vagueness in the environment can be easily handled by fuzzy sets.

**Definition 1.** *Fuzzy Set(FS)* [2]: A fuzzy set (FS)  $X$  on a universe of discourse  $U = (u_1, u_2, \dots, u_n)$  having the membership function  $\mu_X : U \rightarrow [0, 1]$  as follows:

$$X = \{ \langle a, \mu_X(a) \rangle / a \in U \}$$

The membership value  $\mu_X(a)$  describes the degree of the belongingness of  $a \in U$  in  $X$ . When  $\mu_X(a)$  is valued in  $\{0, 1\}$ , it is the characteristic function of a crisp (i.e., non-fuzzy) set.

### Corresponding Author:

Anshu Ohlan, Assistant Professor, Department of Mathematics, All India Jat Heroes Memorial College, Rohtak Haryana, India;  
E-mail: anshu.gahlawat@yahoo.com

Received: 19.02.2017

Revised: 28.02.2017

Accepted: 10.03.2017

The fuzzy divergence measure can be defined as the difference between two fuzzy sets.

Bhandari and Pal [24] established the fuzzy divergence measure analogous to Kullback and Leibler [25] divergence measure, as

$$I(A:B) = \sum_{i=1}^n \left[ \mu_A(x_i) \log \frac{\mu_A(x_i)}{\mu_B(x_i)} + (1-\mu_A(x_i)) \log \frac{1-\mu_A(x_i)}{1-\mu_B(x_i)} \right] \quad (1)$$

with the conditions:

$$I(A:B) = 0 \text{ if } A = B$$

$$I(A:B) = 0 \text{ if } A = B$$

$$I(A:B) \text{ is a convex function of } \mu_A(x_i).$$

## RESULTS

We now propose a generalized measure of Hellinger's divergence between two fuzzy sets  $A$  and  $B$  defined in a universe of discourse  $U = \{a_1, a_2, \dots, a_n\}$  having membership values  $\mu_X(a_i), \mu_Y(a_i) \in (0,1)$  corresponding to Taneja [26] generalized Hellinger's discrimination measure given by

$$h_t(X,Y) = \sum_{i=1}^n \left[ \frac{(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2(t+1)}}{(\sqrt{\mu_X(a_i)\mu_Y(a_i)})^t} + \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2(t+1)}}{(\sqrt{(1-\mu_X(a_i))(1-\mu_Y(a_i))})^t} \right], t \in N \quad (2)$$

**Theorem 1**  $h_t(X,Y)$  is the valid divergence measure of fuzzy sets.

**Proof:** It is clear from (2) that

$$(i) \quad h_t(X,Y) \geq 0$$

$$(ii) \quad h_t(X,Y) = 0 \text{ if } \mu_X = \mu_Y$$

$$(iii) \text{ We now check the convexity of } h_t(X,Y).$$

$$\begin{aligned} & \frac{\partial h_t(X,Y)}{\partial \mu_X(a_i)} \\ &= \frac{(2t+2)(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2(t+1)}}{(\mu_X(a_i)\mu_Y(a_i))^{t/2}} - \frac{1}{2\sqrt{\mu_X(a_i)}} + \left(-\frac{t}{2}\right) \frac{(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2(t+1)}}{(\mu_X(a_i)\mu_Y(a_i))^{\frac{t}{2}-1}} \mu_Y(a_i) \\ &+ 2(t+1) \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2(t+1)}}{(\sqrt{(1-\mu_X(a_i))(1-\mu_Y(a_i))})^t} - \frac{1}{2\sqrt{1-\mu_X(a_i)}} - \frac{t}{2} \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2(t+1)}}{((1-\mu_X(a_i))(1-\mu_Y(a_i)))^{\frac{t}{2}-1}} [-(1-\mu_Y(a_i))] \\ &= (t+1) \left[ \frac{(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2(t+1)}}{\sqrt{\mu_X(a_i)}(\sqrt{\mu_X(a_i)\mu_Y(a_i)})^t} - \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2(t+1)}}{\sqrt{1-\mu_X(a_i)}(\sqrt{(1-\mu_X(a_i))(1-\mu_Y(a_i))})^t} \right] \\ &- \frac{t}{2} \left[ \frac{(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2(t+2)}}{\mu_X(a_i)(\sqrt{\mu_X(a_i)\mu_Y(a_i)})^t} - \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2(t+2)}}{(1-\mu_X(a_i))(\sqrt{(1-\mu_X(a_i))(1-\mu_Y(a_i))})^t} \right] \\ &\frac{\partial^2 h_t(X,Y)}{\partial \mu_X^2(a_i)} = \frac{(t+1)(2t+1)}{2} \left[ \frac{(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2t}}{\mu_X(a_i)(\sqrt{\mu_X(a_i)\mu_Y(a_i)})^t} + \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2t}}{(1-\mu_X(a_i))(\sqrt{(1-\mu_X(a_i))(1-\mu_Y(a_i))})^t} \right] \\ &- \frac{(t+1)(2t+1)}{2} \left[ \frac{(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2t+1}}{\mu_X^{3/2}(a_i)(\sqrt{\mu_X(a_i)\mu_Y(a_i)})^t} + \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2t+1}}{(1-\mu_X(a_i))^{3/2}(\sqrt{(1-\mu_X(a_i))(1-\mu_Y(a_i))})^t} \right] \end{aligned}$$

$$+ \frac{t(t+2)}{4} \left[ \frac{(\sqrt{\mu_X(a_i)} - \sqrt{\mu_Y(a_i)})^{2t+2}}{\mu_X^2(a_i)(\sqrt{\mu_X(a_i)\mu_Y(a_i)})^t} + \frac{(\sqrt{1-\mu_X(a_i)} - \sqrt{1-\mu_Y(a_i)})^{2t+2}}{(1-\mu_X(a_i))^2(\sqrt{(1-\mu_X(a_i))(1-\mu_Y(a_i))})^t} \right] > 0,$$

$$t \in N.$$

Similarly,  $t \in N, t \in N$ .

Thus  $h_t(X,Y)$  is a convex function of fuzzy sets  $X$  and  $Y$  and hence in view of the definition of fuzzy divergence measure of Bhandari and Pal [2] provided in Section 2,  $h_t(X,Y)$  is a valid measure of fuzzy divergence.

**Particular Case:** For  $2h(X,Y)$ ,  $2h(X,Y)$  reduces to  $2h(X,Y)$  where  $h(X,Y)$  is fuzzy Hellinger's divergence measure.

## DISCUSSION

We now provide some more properties of the proposed generalized fuzzy divergence measure (2) in the following theorems. While proving these theorems we consider the distribution of  $U$  in to two parts  $U_1$  and  $U_2$  as

$$U_1 = \{a / a \in U, \mu_X(a_i) \geq \mu_Y(a_i)\} \quad (3)$$

$$\text{and } h_t(X \cup Y, X) + h_t(X \cap Y, X) = h_t(X, Y) \quad (4)$$

### Theorem 2

$$(a) \quad h_t(X \cup Y, X) + h_t(X \cap Y, X) = h_t(X, Y)$$

$$(b) \quad h_t(\overline{X \cup Y}, \overline{X \cap Y}) = h_t(\overline{X} \cap \overline{Y}, \overline{X} \cup \overline{Y}) = h_t(X, Y)$$

$$(c) \quad h_t(X, \overline{X}) = h_t(\overline{X}, X).$$

### Theorem 3

$$(a) \quad h_t(X \cup Y, Z) + h_t(X \cap Y, Z) = h_t(X, Z) + h_t(Y, Z).$$

$$(b) \quad h_t(X, X \cap Y) = h_t(Y, X \cup Y).$$

$$(c) \quad h_t(X, X \cap Y) = h_t(Y, X \cup Y).$$

### Theorem 4

$$(a) \quad h_t(X \cup Y, X \cap Y) = h_t(X, Y).$$

$$(b) \quad h_t(X, \overline{Y}) = h_t(\overline{X}, Y).$$

$$(c) \quad h_t(X, \overline{Y}) = h_t(\overline{X}, Y).$$

$$(d) \quad h_t(X, Y) + h_t(\overline{X}, Y) = h_t(\overline{X}, \overline{Y}) + h_t(X, \overline{Y}).$$

**Proof:** Proof of above theorems follows from (3) and (4).

## CONCLUSION

In the present paper we have obtained generalized information measure of discrimination in fuzzy setting. For this we have proposed and validated the generalized Hellinger's measure of fuzzy divergence. A particular case and a number of the interesting efficient properties of this generalized divergence measure are proven. Finally, we observe that the presence of the parameters in the proposed measure provides a greater flexibility in applications.

## ACKNOWLEDGEMENT

Authors acknowledge the immense help received from the scholars whose article cited and included in references of this manuscript. The authors are also grateful to authors/editors/publishers of all those articles, journals and books from where the literature for this article has been reviewed and discussed.

**Conflict of interest:** Nil

**Source of Funding:** Nil

## REFERENCES

- Shannon, C.E. (1948). A mathematical theory of communication. *The Bell System Technical Journal*, 27(3), 379-423.
- Zadeh, L.A. (1965). Fuzzy sets. *Information and Control*, 8(3), 338-353.
- Zadeh, L.A. (1968). Probability measures of fuzzy events. *Journal of Mathematical Analysis and Applications*, 23(2), 421-427.
- Ohlan A.(2015), "A new generalized fuzzy divergence measure and applications," *Fuzzy Information and Engineering*, 7(4), 507-523.
- Ohlan A.(2016), "Intuitionistic fuzzy exponential divergence: application in multi-attribute decision making," *Journal of Intelligent & Fuzzy Systems*, 30, 1519-1530.
- Ohlan A. and Ohlan R.(2016), "Generalizations of Fuzzy Information Measures," Switzerland: Springer International Publishing.
- Ohlan A. and Ohlan R.(2016), "Fundamentals of Fuzzy Information Measures," in Ohlan A. and Ohlan R., *Generalizations of Fuzzy Information Measures*, Springer International Publishing Switzerland, 1-22.
- Ohlan A. and Ohlan R.(2016), "Parametric Generalized R-norm Fuzzy Information and Divergence Measures," in Ohlan A. and Ohlan R., *Generalizations of Fuzzy Information Measures*, Springer International Publishing Switzerland, 23-52.
- Ohlan A. and Ohlan R.(2016), "Parametric Generalized Exponential Fuzzy Divergence Measure and Strategic Decision-Making," in Ohlan A. and Ohlan R., *Generalizations of Fuzzy Information Measures*, Springer International Publishing Switzerland, 53-69.
- Ohlan A. and Ohlan R.(2016), "Sequence and Application of Inequalities Among Fuzzy Mean Difference Divergence Measures in Pattern Recognition" in Ohlan A. and Ohlan R., *Generalizations of Fuzzy Information Measures*, Springer International Publishing Switzerland, 71-92.
- Ohlan A. and Ohlan R.(2016), "Applications of Generalized Fuzzy Divergence Measure Multi-criteria Decision Making and Pattern Recognition" in Ohlan A. and Ohlan R., *Generalizations of Fuzzy Information Measures*, Springer International Publishing Switzerland, 93-105.
- Ohlan A. and Ohlan R.(2016), "Generalized Hellinger's Divergence Measure and Its Applications" in Ohlan A. and Ohlan R., *Generalizations of Fuzzy Information Measures*, Springer International Publishing Switzerland, 107-121.
- Ohlan A. and Ohlan R.(2016), "Intuitionistic Fuzzy Exponential Divergence and Multi-attribute Decision-Making," in Ohlan A. and Ohlan R., *Generalizations of Fuzzy Information Measures*, Springer International Publishing Switzerland, 123-142.
- Tomar V.P. and Ohlan A. (2014), "Two new parametric generalized R-norm fuzzy information measures", *International Journal of Computer Applications* 93(13), 22-27.
- Tomar V.P. and Ohlan A.(2014), "Sequence of fuzzy divergence measures and inequalities", *AMO - Advanced Modeling and Optimization*, 16(2), 439-452.
- Tomar V.P. and Ohlan A.(2014), "Sequence of inequalities among fuzzy mean difference divergence measures and their applications", *SpringerPlus*, 3, 623, 1-20.
- Tomar V.P. and Ohlan A.(2014), "New parametric generalized exponential fuzzy divergence measure," *Journal of Uncertainty Analysis and Applications*, 2(1), 1-14.
- Ohlan A.(2016), "Some Recent Developments on Probabilistic Information Measures," *International Journal of Innovative Research in Science, Engineering and Technology*, 5(12), 20455-20460.
- Ohlan A.(2016), "Overview on Development of Fuzzy Information Measures," *International Journal of All Research Education and Scientific Methods (IJARESM)*, 4(12), 17-22.
- Ohlan A.(2016), "An Overview on Intuitionistic Fuzzy Similarity Measures," *International Journal of Advanced Technology in Engineering and Science*, 4(11), 192-198.
- Ohlan A.(2016), "Generalized Exponential Fuzzy Information Measures," *International Journal of Advanced Technology in Engineering and Science*, 4(12), 392-399.
- Ohlan A.(2016), "Similarity Measures on Intuitionistic Fuzzy Sets," *International Journal of Science Technology and Management*, 5(12), 463-468.
- Hellinger, E. (1909). Neue begründung der theorie der quadratischen formen von unendlichen vielen veränderlichen. *J. Reine Aug. Math.*, 136, 210-271.
- Bhandari, D., & Pal, N.R. (1993). Some new information measures for fuzzy sets. *Information Sciences*, 67(3), 209-228.
- Kullback, S., & Leibler, R.A. (1951). On information and sufficiency. *The Annals of Mathematical Statistics*, 22(1), 79-86.
- Taneja, I.J. (2013). Seven means, generalized triangular discrimination and generating divergence measures. *Information*, 4, 198-239.